

Author Index

- Abdelghany, A E & Ahmad, M H. Effects of feeding rates on growth and production of Nile tilapia, common carp and silver carp polycultured in fertilized ponds, 415
- Abery, N W, Gunasekera, R M & De Silva, S S. Growth and nutrient utilization of Murray cod *Maccullochella peelii peelii* (Mitchell) fingerlings fed diets with varying levels of soybean meal and blood meal, 279
- Abu-Rezq, T, Al-Abdul-Elah, K, Duremdez, R, Al-Marzouk, A, James, C M, Al-Gharabally, H & Al-Shimmari, J. Studies on the effect of using the rotifer, *Brachionus plicatilis*, treated with different nutritional enrichment media and antibiotics on the growth and survival of blue-fin sea bream, *Sparidentex hasta* (Valenciennes) larvae, 117
- Adams, A see Vatsos, I N
- A-El-Sayed, F M. Effects of stocking density and feeding levels on growth and feed efficiency of Nile tilapia (*Oreochromis niloticus* L.) fry, 621
- Aguayo, S see Roset, J
- Ahmad, M H see Abdelghany, A E
- Ahmed, M see Ayub, Z
- Ahmed, M see Siddiqui, G
- Al-Abdul-Elah, K see Abu-Rezq, T
- Alam, M S see Teshima, S
- Alexis, M see Rigos, G
- Al-Gharabally, H see Abu-Rezq, T
- bin Ali, A see Chong, A
- Allan, G L see Booth, M A
- Al-Marzouk, A see Abu-Rezq, T
- Alongi, D M see McKinnon, A D
- Al-Shimmari, J see Abu-Rezq, T
- Álvarez, M see Roset, J
- Andreassen, B see Jobling, M
- Andrew, J E, Noble, C, Kadri, S, Jewell, H & Huntingford, F A. The effect of demand feeding on swimming speed and feeding responses in Atlantic salmon *Salmo salar* L., gilthead sea bream *Sparus aurata* L. and European sea bass *Dicentrarchus labrax* L. in sea cages, 501
- Andriopoulou, A see Rigos, G
- Anraku, K see Yacoub, S Y
- Archdale, M V see Yacoub, S Y
- Arndt, R E, Routledge, M D, Wagner, E J & Mellenthin, R F. The use of AquaMats® to enhance growth and improve fin condition among raceway cultured rainbow trout *Oncorhynchus mykiss* (Walbaum), 359
- Arnesen, A M see Bendiksen, E Å
- Atse, C B, Audet, C & de la Noüe, J. Effects of temperature and salinity on the reproductive success of Arctic charr, *Salvelinus alpinus* (L.): egg composition, milt characteristics and fry survival, 299
- Audet, C see Atse, C B
- Aupérin, B see Boujard, T
- Ayub, Z & Ahmed, M. A description of the ovarian development stages of penaeid shrimps from the coast of Pakistan, 767
- Baghurst, B C & Mitchell, J G. Sex-specific growth and condition of the Pacific oyster (*Crassostrea gigas* Thunberg), 1253
- Bai, S C see Kim, K H
- Bai, S C see Kim, K-W
- Bai, S C see Wang, X
- Baras, E & Jobling, M. Dynamics of intracohort cannibalism in cultured fish, 461
- de Barniol, L see Jiménez, R
- Barón S B, Bückle, R F & Espina, S. Environmental factors and sexual differentiation in *Poecilia spheonops* Valenciennes (Pisces: Poeciliidae), 615
- Battaglione, S C see Cleary, J J
- Beames, R M see Maina, J G
- Becker, K see Hossain, M A
- Bendiksen, E Å, Jobling, M & Arnesen, A M. Feed intake of Atlantic salmon parr *Salmo salar* L. in relation to temperature and feed composition, 525
- Bhakta, J N & Jana, B B. Influence of sediment phosphorus on utilization efficiency of phosphate fertilizer: a mesocosm study, 203
- Bhatnagar, A see Garg, S K
- Bialowas, H see Brzuska, E
- Blanc, J M. Interaction between diet and genetic aptitude for weight and growth in juvenile rainbow trout, *Oncorhynchus mykiss* (Walbaum), 563
- Blier, P U see Le François, N R
- Bombeo, R F, Fermin, A C & Tan-Fermin, J D. Nursery rearing of the Asian catfish, *Clarias macrocephalus* (Günther), at different stocking densities in cages suspended in tanks and ponds, 1031
- Bombeo-Tuburan, I, Coniza, E B & Rodriguez, E. Preliminary report on nursery and grow-out culture of hatchery-bred grouper (*Epinephelus coioides* Hamilton) in ponds, 379
- Booth, M A, Allan, G L, Evans, A J & Gleeson, V P. Effects of steam pelleting or extrusion on digestibility and performance of silver perch *Bidyanus bidyanus*, 1163
- Borgatti, A R see Pirini, M
- Boujard, T, Labbé, L & Aupérin, B. Feeding behaviour, energy expenditure and growth of rainbow trout in relation to stocking density and food accessibility, 1233
- Boyce, D L see Purchase, C F
- Bray, W A see Perez-Velazquez, M
- Brown, J A see Purchase, C F
- Bruce, M P see Sheppard, J K
- Brzuska, E & Bialowas, H. Artificial spawning of carp, *Cyprinus carpio* (L.), 753
- Bückle, R F see Barón S, B
- Bückle, L F see Hernández, M
- Cameron, C, Gurure, R, Reddy, K, Moccia, R & Leatherland, J F. Correlation between dietary lipid:protein ratios and plasma growth and thyroid hormone levels in juvenile Arctic charr, *Salvelinus alpinus* (Linnaeus), 383
- Campaña-Torres, A see Martínez-Cordova, L R
- Carballo, E C see Ruane, N M
- Caruso, D, Schlumberger, O, Dahm, C & Proteau, J-P. Plasma lysozyme levels in sheatfish *Silurus glanis* (L.) subjected to stress and experimental infection with *Edwardsiella tarda*, 999
- Ceapa, C see Williot, P
- Cecchini, S & Saroglia, M. Antibody response in sea bass (*Dicentrarchus labrax* L.) in relation to water temperature and oxygenation, 607
- Chaitanawisuti, N, Kritsanapuntu, S & Natsukari, Y. Economic analysis of a pilot commercial production for spotted babylon, *Babylonia areolata* (Link 1807), of marketable sizes using a flow-through culture system in Thailand, 1265

- Chan, J C K, Mann, J, Skura, B J, Rowshandeli, M, Rowshandeli, N & Higgs, D A. Effects of feeding diets containing various dietary protein and lipid ratios on the growth performance and pigmentation of post-juvenile coho salmon *Oncorhynchus kisutch* reared in sea water, 1137
- Chen, S-C see Kono, T
- Chen, W-M, Naruse, M & Tabata, M. Circadian rhythms and individual variability of self-feeding activity in groups of rainbow trout *Oncorhynchus mykiss* (Walbaum), 491
- Chereguini, O, García de la Banda, I, Rasines, I & Fernández A. Growth and survival of young turbot (*Scophthalmus maximus* L.) produced with cryopreserved sperm, 637
- Cho, Y L see Lim, L C
- Choi, S-H see Kim, K-W
- Choi, S-M see Kim, K-W
- Choi, Y-H see Kim, K-W
- Chong, A, Hashim, R, Lee, L-C & bin Ali, A. Characterization of protease activity in developing discus *Symphysodon aequifasciata* larva, 663
- Choubert, G see Gouveia, L
- Cleary, J J, Battaglene, S C & Pankhurst, N W. Capture and handling stress affects the endocrine and ovulatory response to exogenous hormone treatment in snapper, *Pagrus auratus* (Bloch & Schneider), 829
- Coloso, R M see Eusebio, P S
- Coniza, E B see Bombeo-Tuburan, I
- Cook, M A & Rust, M B. The effect of light intensity on development and hatching success of lingcod, *Ophiodon elongatus* (Girard), eggs, 217
- Copeman, L A & Parrish, C C. Lipid composition of malpigmented and normally pigmented newly settled yellowtail flounder, *Limanda ferruginea* (Storer), 1209
- Cotter, D V, O'Donovan, Drumm, A, Roche, N, Ling, E N & Wilkins, N P. Comparison of freshwater and marine performances of all-female diploid and triploid Atlantic salmon (*Salmo salar* L.), 43
- Crespo, S see Tort, L
- Cross, T F see Ruff, N
- Cross, T F see Stefánsson, M Ó
- Cui, Y see Wu, L
- Dahm, C see Caruso, D
- Davidson, A see McKinnon, A D
- Davies, S J see White, D A
- De la Roche, J P, Marin, B, Freitas, L & Vélez, A. Embryonic development and larval and post-larval growth of the tropical scallop *Nodipecten (=Lyropecten) nodosus* (L. 1758) (Mollusca: Pectinidae), 819
- De Silva, S S see Abery, N W
- Degani, G. Availability of protein and energy from three protein sources in hybrid sturgeon *Acipenser guldensadtii* x *A. besteri*, 725
- Dennison, W C see Jones, A B
- Dhert, P see Lim, L C
- Divakaran, S see Obaldo, L G
- Dominguez, L see Roset, J
- Downing, G. Impact of spectral composition on larval haddock, *Melanogrammus aeglefinus* L., growth and survival, 251
- Drumm, A see Cotter, D
- Duremdex, R see Abu-Rezq, T
- Ebisuzaki, S see Tomaru, Y
- Ekli, M see Johansen, S J S
- Empis, J see Gouveia, L
- Epp, M A, Ziemann, D A & Schell, D M. Carbon and nitrogen dynamics in zero-water exchange shrimp culture as indicated by stable isotope tracers, 839
- Espe, M see Kvåle, A
- Espina, S see Barón S, B
- Espina, S see Hernández, M
- Eusebio, P S & Coloso, R M. Proteolytic enzyme activity of juvenile Asian sea bass, *Lates calcarifer* (Bloch), is increased with protein intake, 569
- Evans, A J see Booth, M A
- Evensen, T H see Foss, A
- Ezquerro, J M see Ocampo, L
- Falk-Petersen, I-B see Hansen, T K
- Favaloro, E, Lopiano, L & Mazzola, A. Rearing of sharpnout seabream (*Diplodus puntazzo*, Cetti 1777) in a Mediterranean fish farm: monoculture versus polyculture, 137
- Fermin, A C see Bombeo, R F
- Fermin, A C. Effects of alternate starvation and refeeding cycles on food consumption and compensatory growth of abalone, *Haliotis asinina* (Linnaeus), 197
- Fernández, A see Chereguini, O
- Fernández-Garayzabal, J F see Roset, J
- FitzGerald, R D see Ruff, N
- FitzGerald, R D see Stefánsson, M Ó
- Focken, U see Hossain, M A
- Foss, A & Imsland, A K. Compensatory growth in the spotted wolffish *Anarhichas minor* (Olafsen) after a period of limited oxygen supply, 1097
- Foss, A, Evensen, T H & Oiestad, V. Effects of hypoxia and hyperoxia on growth and food conversion efficiency in the spotted wolffish *Anarhichas minor* (Olafsen), 437
- Freites, L see De la Roche, J P
- Furuita, H see Yamamoto, T
- Gallardo, P P, Pedroza-Islas, R, García-Galano, T, Pascual, C, Rosal, C, Sánchez, A & Gaxiola, G. Replacement of live food with microbound diet in feeding *Litopenaeus setiferus* (Burkenroad) larvae, 681
- Gangadhara, B see Nandeesh, M C
- García de la Banda, I see Chereguini, O
- García-Galano, T see Gallardo, P P
- Garg, S K, Kalla, A & Bhatnagar, A. Evaluation of raw and hydrothermally processed leguminous seeds as supplementary feed for the growth of two Indian major carp species, 151
- Gatlin, D M see Perez-Velazquez, M
- Gauthier, R see Michielsens, C G J
- Gaxiola, G see Gallardo, P P
- Gibello, A see Roset, J
- Gleeson, V P see Booth, M A
- Gomes, E see Gouveia, L
- González, M L, Pérez, M C & López, D A. Breeding cycle of the northern scallop, *Argopecten purpuratus* (Lamarck, 1819) in southern Chile, 847
- González-Félix, M L see Perez-Velazquez, M
- Gooley, G J see Ingram, B A
- Gouveia, L, Choubert, G, Pereira, N, Santinha, J, Empis, J & Gomes, E. Pigmentation of gilthead seabream, *Sparus aurata* (L. 1875), using *Chlorella vulgaris* (Chlorophyta, Volvocales) microalgae, 987
- Grant, J see Stewart, A R J
- Grant, W E see Montoya, R A
- Green, B W, Nagdy, Z El & Hebicha, H. Evaluation of Nile tilapia pond management strategies in Egypt, 1037
- Gulyas, T see Williot, P
- Gunasekera, R M see Abery, N W
- Gurure, R see Cameron, C
- Hamre, K see Kvåle, A
- Handley, S J. Optimizing intertidal Pacific oyster (*Thunberg*) culture, Houhora Harbour, northern New Zealand, 1019

- Hansen, T K & Falk-Petersen, I-B. Growth and survival of first-feeding spotted wolffish (*Anarhichas minor* Olafsen) at various temperature regimes, 1119
- Harboe, T see Kvåle, A
- Hart, M J see Penney, R W
- Harvey, S C, Powell, S F, Kennedy, D D, McAndrew, B J & Penman, D J. Karyotype analysis of *Oreochromis mortimeri* (Trewavas) and *Sarotherodon melanotheron* (Rüppell), 339
- Hashim, R see Chong, A
- Hebicha, H see Green, B W
- Hernández, M, Bückle, L F & Espina, S. Temperature preference and acclimation in *Poecilia sphenops* (Pisces, Poeciliidae), 933
- Hickman, R W, Redfearn, P & Tait, M J. Novel effects of salinity and water reuse on growth of juvenile New Zealand turbot, *Colistium nudipinnis* (Waite 1910), a potential aquaculture species, 1009
- Higgs, D A see Chan, J C K
- Higgs, D see Maina, J G
- Hoang, T, Lee, S Y, Keenan, C P & Marsden, G E. Cold tolerance of the banana prawn *Penaeus merguensis* de Man and its growth at different temperatures, 21
- Hoang, T, Lee, S Y, Keenan, C P & Marsden, G E. Spawning behaviour of *Penaeus (Femneropenaeus) merguensis* de Man and the effect of light intensity on spawning, 351
- Hoang, T, Lee, S Y, Keenan, C P & Marsden, G E. Observations on growth, sexual maturity and spawning performance of pond-reared *Penaeus merguensis*, 863
- Hoang, T, Lee, S Y, Keenan, C P & Marsden, G E. Maturation and spawning performance of pond-reared *Penaeus merguensis* in different combinations of temperature, light intensity and photoperiod, 1243
- Hossain, M A, Focken, U & Becker, K. Nutritional evaluation of dhaincha (*Sesbania aculeata*) seeds as dietary protein source for tilapia *Oreochromis niloticus*, 653
- Hotos, G N. Selectivity of the rotifer *Brachionus plicatilis* fed mixtures of algal species with various cell volumes and cell densities, 949
- Huntingford, F A see Andrew, J E
- Husnah & Lin, C K. Toxicity of chlorine to different sizes of black tiger shrimp (*Penaeus monodon*) in low-salinity shrimp pond water, 1129
- Hwang, Y J see Kim, K H
- Ibrahim, C-O see Ng, W-K
- Imsland, A K see Foss, A
- Ingram, B A, McKinnon, L J & Gooley, G J. Growth and survival of selected aquatic animals in two saline groundwater evaporation basins: an Australian case study, 425
- Ishikawa, M see Teshima, S
- Iwama, G see Maina, J G
- Jafri, A K see Hotos, G N
- James, C M see Abu-Rezq, T
- Jana, B B see Bhakta, J N
- Jeffs, A G see Sheppard, J K
- Jensen, L see Ringo, E
- Jerry, D R, Purvis, I W & Piper, L R. Genetic differences in growth among wild populations of the yabby, *Cherax destructor* (Clark), 917
- Jewell, H see Andrew, J E
- Jiménez, R, de Barniol, L & Machuca, M. *Nematopsis marinus* n. sp., a new septate gregarine from cultured penaeoid shrimp *Litopenaeus vannamei* (Boone), in Ecuador, 231
- Jobling, M see Baras, E
- Jobling, M see Bendiksen, E Å
- Jobling, M see Johansen, S J S
- Jobling, M. Editorial, 459
- Jobling, M, Andreassen, B, Larsen, A V & Olsen, R L. Fat dynamics of Atlantic salmon *Salmo salar* L. smolt during early seawater growth, 739
- Jobling, M, Larsen, A V, Andreassen, B & Olsen, R L. Adiposity and growth of post-smolt Atlantic salmon *Salmo salar* L., 533
- Jobling, M, Larsen, A V, Andreassen, B, Sigholt, T & Olsen, R L. Influence of a dietary shift on temporal changes in fat deposition and fatty acid composition of Atlantic salmon post-smolt during the early phase of seawater rearing, 875
- Johansen, S J S, Ekli, M & Jobling, M. Is there lipostatic regulation of feed intake in Atlantic salmon *Salmo salar* L.? 515
- Johnston, D, Lourey, M, Van Tien, D, Luu, T T & Xuan, T T. Water quality and plankton densities in mixed shrimp-mangrove forestry farming systems in Vietnam, 785
- Jones, A B, Preston, N P & Dennison, W C. The efficiency and condition of oysters and macroalgae used as biological filters of shrimp pond effluent, 1
- Jones, D A see Puello-Cruz, A C
- Jones, P L, Thanuthong, T & Kerr, P. Preliminary study on the use of synthetic substrate for juvenile stage production of the yabby, *Cherax destructor* (Clark) (Decapoda: Parastacidae), 811
- Kadri, S see Andrew, J E
- Kalla, A see Garg, S K
- Kanazawa, A see Teshima, S
- Karl, H, Kuhlmann, H & Oetjen, K. Transfer of toxaphene and chlordane into farmed rainbow trout, *Oncorhynchus mykiss* (Walbaum) via feed, 925
- Kawabata, Z see Tomaru, Y
- Keenan, C P see Hoang, T
- Kelly, M S & Owen, P V. Growth of the abalone *Haliotis tuberculata* L. at Scottish sea temperatures, 729
- Kennedy, D D see Harvey, S C
- Kentouri, M see Paspatis, M
- Kerr, P see Jones, P L
- Kerry, J P see Ruff, N
- Kim, D S see Kim, K H
- Kim, K see Wang, X
- Kim, K H, Hwang, Y J, Kim, K W, Bai, S C & Kim, D S. Effects of dietary aloe on chemiluminescent responses of peripheral blood phagocytes and resistance against *Edwardsiella tarda* Ewing and McWhorter 1965, in the cultured olive flounder, *Paralichthys olivaceus* (Temminck et Schlegel), 147
- Kim, K W see Kim, K H
- Kim, K W, Wang, X J & Bai, S C. Optimum dietary protein level for maximum growth of juvenile olive flounder *Paralichthys olivaceus* (Temminck et Schlegel), 673
- Kim, K-W, Wang, X, Choi, S-M, Bai, S C, Choi, Y-H & Choi, S-H. Long-term evaluation of extruded diets compared with raw fish moist diet for growing Korean rockfish, *Sebastes schlegelii* (Hilgendorf), 979
- Kisia, S M see Maina, J G
- Kiyohara, S see Yacoub, S Y
- Kloosterboer, R J see Lambooi, E
- Komen, J see Ruane, N M
- Kono, T, Okamoto, K & Sakai, M. Analysis of expressed sequence tags from a yellowtail, *Seriola quinqueradiata* (Temminck & Schlegel), head kidney cDNA library, 291
- Kono, T, Ooyama, T, Chen, S-C & Sakai, M. Sequencing of 16S-23S rRNA internal transcribed spacer and its application in the identification of *Nocardia seriolae* by polymerase chain reaction, 1195
- Koshio, S see Teshima, S
- Kritsanapuntu, S see Chaitanawisuti, N
- Kuhlmann, H see Karl, H
- Kuhlmann, H see Lambooi, E

- Kumatabara, Y see Tomaru, Y
- Kvåle, A, Harboe, T, Espe, M, Næss, T & Hamre, K. Effect of predigested protein on growth and survival of Atlantic halibut larvae (*Hippoglossus hippoglossus* L.), 311
- Kwan, T S see Yusoff, F M
- Labbé, L see Boujard, T
- Lambooi, E, van de Vis, J W, Kloosterboer, R J & Pieterse, C. Evaluation of head-only and head-to-tail electrical stunning of farmed eels (*Anguilla anguilla*, L.) for the development of a humane slaughter method, 323
- Lambooi, E, van de Vis, J W, Kuhlmann, H, Münkner, W, Oehlenschläger, J, Kloosterboer, R J & Pieterse, C. A feasible method for humane slaughter of eel (*Anguilla anguilla* L.): electrical stunning in fresh water prior to gutting, 643
- Láron, M A see Santiago, C B
- Larsen, A V see Jobling, M
- Lawrence, A L see Montoya, R A
- Lawrence, A L see Perez-Velazquez, M
- Le François, N R, Lemieux, H & Blier, P U. Biological and technical evaluation of the potential of marine and anadromous fish species for cold-water mariculture, 95
- Le Vay, L see Puello-Cruz, A C
- Leatherland, J F see Cameron, C
- Lee, L-C see Chong, A
- Lee, S Y see Hoang, T
- Lemieux, H see Le François, N R
- Lim, H-A see Ng, W-K
- Lim, L C, Cho, Y L, Dhert, P, Wong, C C, Nelis, H & Sorgeloos, P. Use of decapsulated *Artemia* cysts in ornamental fish culture, 575
- Lim, S-L see Ng, W-K
- Lin, C K see Husnah
- Lindström-Seppä, P see Ritola, O
- Ling, E N see Cotter, D
- Livingstone, D R see Ritola, O
- Lodemel, J B see Ringo, E
- López, D A see González, M L
- Lopiano, L see Favalaro, E
- Lorenzen, K see Michielsens, C G J
- Lourey, M see Johnston, D
- Lund, V see Ringo, E
- Luu, T T see Johnston, D
- McAndrew, B J see Harvey, S C
- Machuca, M see Jiménez, R
- McKinnon, A D, Trott, L A, Alongi, D M & Davidson, A. Water column production and nutrient characteristics in mangrove creeks receiving shrimp farm effluent, 55
- McKinnon, L J see Ingram, B A
- McLeod, I, Pantus, F & Preston, N. The use of a geographical information system for land-based aquaculture planning, 241
- Madhusoodana Kurup, B see Ranjeet, K
- Maina, J G, Beames, R M, Higgs, D, Mbugua, P N, Iwama, G & Kisia, S M. Digestibility and feeding value of some feed ingredients fed to tilapia *Oreochromis niloticus* (L.), 853
- Manissery, J K see Nandeesha, M C
- Mann, J see Chan, J C K
- Maragoudaki, D see Paspatis, M
- Marin, B see De la Roche, J P
- Marinho-Soriano, E, Morales, C & Moreira, C W S. Cultivation of *Gracilaria* (Rhodophyta) in shrimp pond effluents in Brazil, 1081
- Marsden, G E see Hoang, T
- Martinez-Cordova, L R, Campaña-Torres, A & Porchas-Cornejo, M A. Promotion and contribution of biota in low water exchange ponds farming blue shrimp *Litopenaeus stylirostris* (Stimpson), 27
- Martinez-Cordova, L R, Campaña-Torres, A & Porchas-Cornejo, M A. The effects of variation in feed protein level on the culture of white shrimp, *Litopenaeus vannamei* (Boone) in low-water exchange experimental ponds, 995
- Martinez-Palacios, C A see Olvera-Novoa, M A
- Matias, H B see Yusoff, F M
- Matsuoka, T see Yacoub, S Y
- Mayhew, T M see Ringo, E
- Mazzola, A see Favalaro, E
- Mbugua, P N see Maina, J G
- Medina, A see Nieves, M
- Mellenthin, R F see Arndt, R E
- Michielsens, J C G, Lorenzen, K, Phillips, M J & Gauthier, R. Asian carp farming systems: towards a typology and increased resource use efficiency, 403
- Milstein, A see Wahab, M A
- Milstein, A, Wahab, M A & Rahman, M M. Environmental effects of common carp *Cyprinus carpio* (L.) and mrigal *Cirrhinus mrigala* (Hamilton) as bottom feeders in major Indian carp polycultures, 1103
- Mitchell, J G see Baghurst, B C
- Moccia, R see Cameron, C
- Montoya, R A, Lawrence, A L, Grant, W E & Velasco, M. Simulation of inorganic nitrogen dynamics and shrimp survival in an intensive shrimp culture system, 81
- Moody, A J see White, D A
- Morales, C see Marinho-Soriano, E
- Moreira, W S C see Marinho-Soriano, E
- Münkner, W see Lambooi, E
- Muñoz, M J see Roset, J
- Myklebust, R see Ringo, E
- Næss, T see Kvåle, A
- Nagdy, Z El see Green, B W
- Nakano, S see Tomaru, Y
- Nakano, S-I see Tomaru, Y
- Nam, C Q see Vromant, N
- Nandeesha, M C, Gangadhara, B & Manissery, J K. Further studies on the use of mixed feeding schedules with plant- and animal-based diets for common carp *Cyprinus carpio* (Linnaeus), 1157
- Naruse, M see Chen, W-M
- Natsukari, Y see Chaitanawisuti, N
- Nelis, H see Lim, L C
- Nengas, I see Rigos, G
- Ng, W-K, Lim H-A, Lim, S-L & Ibrahim, C-O. Nutritive value of palm kernel meal pretreated with enzyme or fermented with *Trichoderma koningii* (Oudemans) as a dietary ingredient for red hybrid tilapia (*Oreochromis* sp.), 1199
- Nieves, M, Voltolina, D, Medina, A, Pia, P & Ruiz, J L. Zeolites and diatom growth, 75
- Noble, C see Andrew, J E
- de la Noüe, J see Atse, C B
- O'Donovan, V see Cotter, D
- Obaldo, L G, Divakaran, S & Tacon, A G. Method for determining the physical stability of shrimp feeds in water, 369
- Ocampo, L & Ezquerro, J M. Digestive protease activity in juvenile *Farfantepenaeus californiensis* as a function of dissolved oxygen and temperature, 1073
- Oehlenschläger, J see Lambooi, E
- Oetjen, K see Karl, H
- Oiestad, V see Foss, A
- Okamoto, K see Kono, T
- Oliva-Teles, A see Pereira, T G

- Olivera-Castillo, L see Olivera-Novoa, M A
 Ollevier, F see Vromant, N
 Olsen, R E see Ringo, E
 Olsen, R L see Jobling, M
 Olivera-Novoa, M A, Olivera-Castillo, L & Martínez-Palacios, C A. Sunflower seed meal as a protein source in diets for *Tilapia rendalli* (Boulanger, 1896) fingerlings, 223
 Ooyama, T see Kono, T
 Owen, P V see Kelly, M S
 Ozbay, G & Riley, J G. An analysis of refractometry as a method of determining blood total protein concentration in the American lobster *Homarus americanus* (Milne Edwards), 557
 Padrós, F see Tort, L
 Page, G I see White, D A
 Pagliarini, A see Pirini, M
 Pankhurst, N W see Cleary, J J
 Pantus, F see McLeod, I
 Parra, G & Yufera, M. Tolerance response to water pH in larvae of two marine fish species, gilthead seabream, *Sparus aurata* (L.) and Senegal sole, *Solea senegalensis* (Kaup), during development, 747
 Parrish, C C see Copeman, L A
 Pascual, C see Gallardo, P P
 Paspatis, M, Maragoudaki, D & Kentouri, M. Feed discrimination and selection in self-fed European sea bass *Dicentrarchus labrax* L., 509
 Pease, B C see Walsh, C T
 Pedroza-Islas, R see Gallardo, P P
 Penman, D J see Harvey, S C
 Penney, R W, Hart, M J & Templeman, N. Comparative growth of cultured blue mussels, *Mytilus edulis*, *M. trossulus* and their hybrids, in naturally occurring mixed-species stocks, 693
 Pereira, N see Gouveia, L
 Pereira, T G & Oliva-Teles, A. Preliminary evaluation of pea seed meal in diets for gilthead sea bream (*Sparus aurata*) juveniles, 1183
 Pérez, M C see González, M L
 Perez-Velazquez, M, Lawrence, A L, Gatlin, D M, III, González-Félix, M L & Bray, W A. Replacement of fresh dietary components by a dry feed for successful maturation of male *Litopenaeus vannamei* (Boone) broodstock, 1091
 Peters, L D see Ritola, O
 Petrusdottir, T E. Influence of feeding frequency on growth and size dispersion in Arctic charr *Salvelinus alpinus* (L.), 543
 Phillips, M J see Michielsens, C G J
 Pia, P see Nieves, M
 Pieterse, C see Lambooi, E
 Pieterse, C see Lambooi, E
 Piper, L R see Jerry, D R
 Pirini, M, Pagliarini, A, Ventrella, V, Trombetti, F, Trigari, G & Borgatti, A R. Response to T₃ treatment and changing environmental salinity of liver lipid composition, mitochondrial respiration and (Na⁺ + K⁺)-ATPase activity in rainbow trout *Oncorhynchus mykiss* Walbaum, 891
 Porchas-Cornejo, M A see Martínez-Cordova, L R
 Powell, S F see Harvey, S C
 Preston, N P see Jones, A B
 Preston, N see McLeod, I
 Proteau, J-P see Caruso, D
 Puello-Cruz, A C, Sangha, R S, Jones, D A & Le Vay, L. Trypsin enzyme activity during larval development of *Litopenaeus vannamei* (Boone) fed on live feeds, 333
 Puigcerver, M see Tort, L
 Purchase, C F, Boyce, D L & Brown, J A. Occurrence of hypome-lanization in cultured yellowtail flounder *Limanda ferruginea*, 1191
 Purvis, I W see Jerry, D R
 Rahman, M M see Milstein, A
 Rahman, M M see Wahab, M A
 Ranjeet, K & Madhusoodana Kurup, B. Management strategies associating batch-graded and size-graded postlarvae can reduce heterogeneous individual growth in *Macrobrachium rosenbergii* (de Man), 1221
 Rasines, I see Chereguini, O
 Reddy, K see Cameron, C
 Redfearn, P see Hickman, R W
 Rigos, G, Alexis, M, Andriopoulou, A & Nengas, I. Temperature-dependent pharmacokinetics and tissue distribution of oxolinic acid in sea bass, *Dicentrarchus labrax* L., after a single intravascular injection, 1175
 Riley, J G see Ozbay, G
 Ringo, E, Lodemel, J B, Myklebust, R, Jensen, L, Lund, V, Mayhew, T M & Olsen, R E. The effects of soybean, linseed and marine oils on aerobic gut microbiota of Arctic charr *Salvelinus alpinus* L. before and after challenge with *Aeromonas salmonicida* ssp. *Salmonicida*, 591
 Ritola, O, Peters, L D, Livingstone, D R & Lindström-Seppä, P. Effects of *in vitro* exposure to ozone and/or hyperoxia on superoxide dismutase, catalase, glutathione and lipid peroxidation in red blood cells and plasma of rainbow trout, *Oncorhynchus mykiss* (Walbaum), 165
 Robb, F, D H, Wotton, S B & van de Vis, J W. Preslaughter electrical stunning of eels, 37
 Roche, N see Cotter, D
 Rodriguez, E see Bombeo-Tuburan, I
 Rosal, C see Gallardo, P P
 Roset, J, Gibello, A, Aguayo, S, Domínguez, L, Álvarez, M, Fernández-Garayzabal, J F, Zapata, A & Muñoz, M J. Mortality of rainbow trout [*Oncorhynchus mykiss* (Walbaum)] associated with freshwater dinoflagellate bloom [*Peridinium polonicum* (Woloszynska)] in a fish farm, 141
 Routledge, M D see Arndt, R E
 Rowshandel, M see Chan, J C K
 Rowshandel, N see Chan, J C K
 Ruane, N M, Carballo, E C & Komen, J. Increased stocking density influences the acute physiological stress response of common carp *Cyprinus carpio* (L.), 777
 Ruff, N, FitzGerald, R D, Cross, T F & Kerry, J P. Fillet shelf-life of Atlantic halibut *Hippoglossus hippoglossus* L. fed elevated levels of α -tocopheryl acetate, 1059
 Ruff, N, FitzGerald, R D, Cross, T F, Teurtrie, G & Kerry, J P. Slaughtering method and dietary α -tocopheryl acetate supplementation affect *rigor mortis* and fillet shelf-life of turbot (*Scophthalmus maximus* L.), 703
 Ruiz, J L see Nieves, M
 Rust, M B see Cook, M A
 Sakai, M see Kono, T
 Sánchez, A see Gallardo, P P
 Sangha, R S see Puello-Cruz, A C
 Santiago, C B & Laron, M A. Growth and fry production of Nile tilapia, *Oreochromis niloticus* (L.), on different feeding schedules, 129
 Santinha, J see Gouveia, L
 Saroglia, M see Cecchini, S
 Schell, D M see Epp, M A
 Schlumberger, O see Caruso, D
 Sheppard, J K, Bruce, M P & Jeffs, A G. Optimal feed pellet size for culturing juvenile spiny lobster *Jasus edwardsii* (Hutton, 1875) in New Zealand, 913
 Shima, T see Yamamoto, T
 Siddiqui, G & Ahmed, M. Gametogenic patterns of the larviparous oyster *Ostrea nomades* from Karachi, Pakistan (northern Arabian Sea), 1049

- Sigholt, T *see* Jobling, M
 Skura, B J *see* Chan, J C K
 Sorgeloos, P *see* Lim, L C
 Stefánsson, M Ó, Fitzgerald, R D & Cross, T F. Growth, feed utilization and growth heterogeneity in juvenile turbot *Scophthalmus maximus* (Rafinesque) under different photoperiod regimes, 177
 Stewart, A R J & Grant, J. Disaggregation rates of extruded salmon feed pellets: influence of physical and biological variables, 799
 Suzuki, N *see* Yamamoto, T
 Swaile, J *see* White, D A
- Tabata, M *see* Chen, W-M
 Tacon, A G *see* Obaldo, L G
 Tait, M J *see* Hickman, R W
 Tan-Fermin, J D *see* Bomboe, R F
 Templeman, N *see* Penney, R W
 Tendencia, E A. *Vibrio harveyi* isolated from cage-cultured seabass *Lates calcarifer* Bloch in the Philippines, 455
 Teshima, S, Alam, M S, Koshio, S, Ishikawa, M & Kanazawa, A. Assessment of requirement values for essential amino acids in the prawn, *Marsupenaeus saponicus* (Bate), 395
 Teurtrie, G *see* Ruff, N
 Thanuthong, T *see* Jones, P L
 Thompson, K D *see* Vatsos, I N
 Tomaru, Y, Ebisuzaki, S, Kawabata, Z & Nakano, S-I. Respiration rates of the Japanese pearl oyster, *Pinctada fucata martensii*, feeding on *Pavlova lutheri* and *Chaetoceros gracilis*, 33
 Tomaru, Y, Kumatabara, Y, Kawabata, Z & Nakano, S. Effect of water temperature and chlorophyll abundance on shell growth of the Japanese pearl oyster, *Pinctada fucata martensii*, in suspended culture at different depths and sites, 109
 Tort, L, Puigcerver, M, Crespo, S & Padrós, F. Cortisol and haematological response in sea bream and trout subjected to the anaesthetics clove oil and 2-phenoxyethanol, 907
 Trigari, G *see* Pirini, M
 Trombetti, F *see* Pirini, M
 Trott, L A *see* McKinnon, A D
- Ulloa R, J B & Verreth, J J A. Growth, feed utilization and nutrient digestibility in tilapia fingerlings (*Oreochromis aureus* Steindachner) fed diets containing bacteria-treated coffee pulp, 189
 Usmani, N *see* Hotos, G N
 Usmani, N & Jafri, A K. Effect of fish size and temperature on the utilization of different protein sources in two catfish species, 959
- Van Tien, D *see* Johnston, D
 Vatsos, I N, Thompson, K D & Adams, A. Development of an immunofluorescent antibody technique (IFAT) and *in situ* hybridization to detect *Flavobacterium psychrophilum* in water samples, 1087
 Velasco, M *see* Montoya, R A
 Vélez, A *see* De la Roche, J P
 Ventrella, V *see* Pirini, M
 Verreth, J A J *see* Ulloa, R J B
 van de Vis, J W *see* Lambooi, E
- van de Vis, J W. *see* Robb, F
 Voltolina, D *see* Nieves, M
 Vromant, N, Nam, C Q & Ollevier, F. Growth performance and use of natural food by *Oreochromis niloticus* (L.) in polyculture systems with *Barbodes gonionotus* (Bleeker) and *Cyprinus carpio* (L.) in intensively cultivated rice fields, 969
- Wagner, E J *see* Arndt, R E
 Wahab, M A *see* Milstein, A
 Wahab, M A, Rahman, M M & Milstein, A. The effect of common carp, *Cyprinus carpio* (L.) and mrigal, *Cirrhinus mrigala* (Hamilton) as bottom feeders in major Indian carp polycultures, 547
 Walsh, C T & Pease, B C. The use of clove oil as an anaesthetic for the longfinned eel, *Anguilla reinhardtii* (Steindachner), 627
 Wang, J & Xia, D. Studies on fish heterosis with DNA fingerprinting, 941
 Wang, X *see* Kim, K-W
 Wang, X, Kim, K & Bai, S C. Effects of different dietary levels of L-ascorbyl-2-polyphosphate on growth and tissue vitamin C concentrations in juvenile olive flounder, *Paralichthys olivaceus* (Temminck et Schlegel), 261
 Wang, X J *see* Kim, K W
 White, D A, Page, G I, Swaile, J, Moody, A J & Davies, S J. Effect of esterification on the absorption of astaxanthin in rainbow trout, *Oncorhynchus mykiss* (Walbaum), 343
 Wilkins, N P *see* cotter, D
 Williot, P, Gulyas, T & Ceapa, C. An analogue of GnRH is effective for induction of ovulation and spermiation in farmed Siberian sturgeon, *Acipenser baerii* Brandt, 735
 Wong, C C *see* Lim, L C
 Wotton, D H *see* Robb, F
 Wootton, R J *see* Wu, L
 Wu, L, Xie, S, Zhu, X, Cui, Y & Wootton, R J. Feeding dynamics in fish experiencing cycles of feed deprivation: a comparison of four species, 481
- Xia, D *see* Wang, J
 Xie, S *see* Wu, L
 Xuan, T T *see* Johnston, D
- Yacoub, S Y, Anraku, K., Archdale, M V, Matsuoka, T & Kiyohara, S. Exposure of taste buds to potassium permanganate and formalin suppress the gustatory neural response in the Nile tilapia, *Oreochromis niloticus* (Linnaeus), 445
 Yamamoto, T, Shima, T, Furuta, H & Suzuki, N. Influence of dietary fat level and whole-body adiposity on voluntary energy intake by juvenile rainbow trout *Oncorhynchus mykiss* (Walbaum) under self-feeding conditions, 715
 Yüfera, M *see* Parra, G
 Yusoff, F M, Zubaidah, M S, Matias, H B & Kwan, T S. Phytoplankton succession in intensive marine shrimp culture ponds treated with a commercial bacterial product, 269
- Zapata, A *see* Roset, J
 Zhu, X *see* Wu, L
 Ziemann, D A *see* Epp, M A
 Zubaidah, M S *see* Yusoff, F M

Keyword Index

- 16S-23S ITS, 1197
- abalone, 197, 733
- absorption, 343
- Acanthopagrus butcheri*, 425
- Acipenser baerii*, 737
- adaptive trials, 1221
- adiposity, 715
- aerobic gut microbiota, 591
- age, 863
- α -tocopherol, 703, 1059
- alternate feeding, 1157
- alternative protein sources, 1183
- amino acid, 395
- anaesthetic, 627
- anchovy fish meal, 853
- ANFs, 189
- Anguilla*, 627
- antibiotics, 117
- antibody response, 607
- antinutrient factors, 151
- antioxidant
- defence, 1059
 - enzymes, 165
- apparent digestibility coefficient (ADC), 959
- aquaculture, 461, 501, 563, 799, 913
- planning, 241
 - species, 95
- Aquamats®, 359
- Arcopecten purpuratus*, 847
- Arctic charr *Salvelinus alpinus* (L.), 299, 543, 591
- Artemia* replacement, 681
- artificial diet, 733
- artificial spawning, 737, 753
- Asia, 403
- Asian catfish, 1031
- astaxanthin pigmentation, 1137
- astaxanthin, 343
- Asteromonas*, 949
- Atlantic halibut, 311
- Atlantic salmon *Salmo salar* L., 515
- Babylonia areolata*, 1265
- bacterial product, 269
- Barbodes gonionotus* (Bleeker), 969
- batch grading, 1221
- behaviour, 461
- benzocaine, 627
- Bidyanus bidyanus*, 425
- extrusion, 1163
- biochemical indices, 140
- biofiltration, 1
- biomass, 1081
- biota promotion, 27
- bivalves, 819, 1253
- blood meal, 279
- blood protein refractometry, 557
- blue shrimp, 27
- body composition, 515, 533, 739, 875
- Brachionus plicatilis*, 949
- breeding programme, 913
- broken-line model, 673
- cage nursery, 1031
- cannibalism, 461
- carcass proximate analysis, 383
- carotenoids, 987
- carp, 403, 753, 941
- pituitary, 737
- catla, 547
- cDNA, 291
- Chaetoceros*, 75, 949
- challenge disease, 999
- characterization, 663
- Chile, 847
- chlordan, 925
- Chlorella*, 949
- vulgaris* (Chlorophyta, Volvocales), 987
- chlorine, 1129
- chlorophyll, 109
- chymotrypsin, 1073
- circadian rhythm, 491
- Clarias gariepinus* (Burchell, 1822), 959
- Clarias macrocephalus*, 1031
- clove oil, 627
- coffee pulp, 189
- coho salmon, 1137
- cold-water aquaculture, 95
- Colistium nudipinnis*, 1009
- Colocasia, 1157
- colour, 703, 1059
- common carp, 415, 547, 777, 1103
- compensatory growth, 197, 1101
- competition, 969
- condition factor, 999
- condition, 1, 1253
- consumption, 481
- Crassostrea gigas*, 425, 1019, 1253
- cryopreserved sperm, 637
- culture, 819
- Cyprinus carpio* (L.), 969, 1157
- decapsulated *Artemia* cysts, 575
- demand feeding, 501
- development, 663
- DHA Protein Selco, 117
- DHA, 1209
- diagnosis, 1197
- diet size, 913
- diet, 563, 1095
- dietary fat, 533, 715, 739
- dietary levels, 261
- dietary lipid, 1137
- dietary lipid:protein ratios, 383
- dietary oils, 591
- dietary protein, 1137
- digestibility, 279, 725, 1163, 1183, 1199
- digestive enzymes, 333

- digestive tract, 591
Diplodus puntazzo, 140
discus, 663
dissolved oxygen, 1073
DNA fingerprinting, 941
dominance hierarchies, 177
dominance, 491
Dunaliella, 949
Duratón river, 145
- economic analysis, 1265
economics, 1037
Edwardsiella tarda, 999
eel, 37, 323, 627, 643
efficiency, 1
effluents, 1081
egg development, 217
egg, 299
Egypt, 1037
electrical stunning, 323, 643
electricity, 37
electrophysiology, 445
embryo, 217
embryonic development, 819
energy balance, 533
energy intake, 715
Enterogonia orbicularis, 1019
environmental temperature, 607
essential fatty acids, 1209
esters, 343
European sea bass *Dicentrarchus labrax* L., 509
expressed sequence tag, 291
extensive shrimp culture, 785
extruded pellet, 979
- fat storage, 875
fatty acids, 875
feed
 composition, 525
 cost, 129
 discrimination, 509
 enzyme, 1199
 intake, 461, 525
 pellets, 799
 preference, 515
 utilization, 177
feeding and nutrition crustaceans, 681
feeding behaviour, 491, 501
feeding frequency, 543
feeding levels, 621
feeding rate, 415
feeding study, 925
feeding, 811
fertilization, 351
fertilizer dose, 203
filtration, 949
fin erosion, 359
first-feeding, 1119
fish by-catch, 569
fish, 425
 farm, 145
 larvae, 747
 meal substitution, 279
 meal, 725
 nutrition, 189
fishmeal replacer, 979
flatworm, 1019
Flavobacterium psychrophilum, 1090
flow-through seawater system, 1265
flume, 799
food concentration, 33
formalin, 445
formulated diet, 311, 569
fouling, 1
fresh water, 643
freshwater aquaculture, 403
freshwater dinoflagellate, 145
fry survival, 299
- gametogenesis, 1049, 1253
gametogenic cycle, 847
genetic base, 913
genetics, 563
genotype, 693
gibel carp, 481
gilthead seabream, 987, 1183
GIS, 241
glutathione, 165
GnRH α , 737
gonadal steroids, 829
Gracilaria, 1081
gregarine, 231
growing-out, 1265
growth 43, 75, 109, 151, 177, 251, 261, 279, 359, 461, 491, 533, 543, 563, 637, 693, 733, 739, 811, 913, 1009, 1119, 1157, 1193
 compensation, 481
 performance, 140, 1137
 rate, 383, 1081, 1253
guppy, 575
gustatory, 445
- haddock, 251
haematocrit, 999
halibut *Hippoglossus hippoglossus* L., 1059
Haliotis asinina, 197
Haliotis tuberculata, 733
harmful blooms, 145
hermaphrodites, 1049
heterogeneous individual growth (HIG), 1221
Heteropneustes fossilis (Bloch), 959
heterosis, 941
high-fibre sunflower cake, 853
Hippoglossus hippoglossus L., 311
hormone assay validation, 383
hybrids, 693
hydrolysed protein, 311
hyperoxia, 165, 1101
hyperphagia, 481
hypomelanization, 1193
hypoxia, 1101
- IFAT detection, 1090
in situ hybridization, 1090
Indian major carp, 151
indirect ELISA, 607
induced ovulation, 829
ingestion, 949
initial sediment phosphate, 203
integrated farming, 785
integrated systems, 403
intestine, 343
- juvenile olive flounder, 261
juvenile turbot, 177, 1009

- karyotype evolution, 339
 kidney, 291
 Korean rockfish, 979
 larvae, 251, 311, 663
 larval fish, 1209
 larval growth, 819
 larviparous, 1049
 leguminous seeds, 151
 leucocrit, 999
 light intensity, 217, 351, 1243
 light, 251
Limanda ferruginea, 1193, 1209
 lingcod, 217
 lipid deposition, 515
 lipid oxidation, 703, 1059
 lipostat, 515
Litopenaeus
 setiferus larvae, 681
 stylirostris, 27
 vannamei (Boone), 333, 839, 995, 1095
 live shipment, 557
 liver lipids, 891
 lobsters, 557
 longsnout catfish, 481
 low salinity, 1129
 low-fibre sunflower cake, 853
Macquaria novemaculeata, 425
 macroalgae, 1
Macrobrachium rosenbergii, 1221
 mangroves, 785
 mariculture diversification, 95
Marsupenaeus japonicus, 395, 425
 maturity, 863
 metaphase chromosomes, 339
 microalga, 987
 microbound diet, 681
 milt, 299
 minnow, 481
 mitochondrial respiration, 891
 mixed feeding, 1157
 modelling, 81
 moist pellet, 979
 molly, 575
 Mpi, 693
 mrigal, 547, 1103
 mudworm, 1019
 multivariate statistics, 1103
 Murray cod, 279
Mytilus
 edulis, 693
 trossulus, 693
 ($\text{Na}^+ + \text{K}^+$)-ATPase, 891
 natural food, 27
Nematopsis marinus n.sp., 231
 New Zealand, 1009
 Nile tilapia, 415, 621, 1037
 nitrogen dynamics, 81
 nocardiosis, 1197
Nodipecten (= *Lyropecten*) *nodosus*, 891
 normoxia, 1101
 nutrient leaching, 369
 nutrition, 839
 olive flounder, 673
 omena fishmeal, 853
Oncorhynchus mykiss (Walbaum), 165, 425, 715
 ontogeny, 461
Oreochromis, 189
 niloticus (L.), 969
Ostrea nomades, 1049
 ovarian
 maturation, 1243
 stages, 767
 Ovopel, 753
 oxidative stress, 165
 oxolinic acid, 1175
 oyster, 1, 425, 1019, 1049
 ozone, 165
Pagrus auratis, 425
 palm kernel meal, 1199
Paralichthys olivaceus, 673
 parasite, 231
 particulates, 1, 799
 PCR, 1197
 pea seed meal, 1183
 pearl oyster, 33, 109
 pellet stability, 369
 penaeid shrimps, 767
Penaeus
 (*Fenneropenaeus*) *merguensis*, 351
 merguensis, 863, 1243
 merguensis and growth, 21
 monodon, 425
Peridinium polonicum, 145
 pH, 747
 pharmacokinetics, 1175
 phosphorus dynamics, 203
 photoperiod, 1243
 manipulation, 177
 phytoplankton succession, 269
 pigmentation, 987, 1193, 1209
 pituitary, 753
 plant protein, 223
 plasma
 cortisol, 777
 lysozyme, 999
 platy, 575
Pleuronectes ferrugineus, 1193
Proecilia sphenops (Pisces Peociliidae), 933
 polyculture, 415, 547, 969, 1103
Polydora hoplura, 1019
 pond-reared broodstock, 863
 potassium permanganate, 445
 poultry meal, 725
 practical diets, 129
 prawn, 395
 preference, 509
 preferred and avoided temperature, 933
 production systems, 1037
 protease, 663
 protein, 569, 959
 level, 995
 requirement, 673
 protein:lipid ratio, 1137
 proteolytic enzyme, 569
 punti, 547, 1103
 Rainbow trout fry syndrome, 1090
 rainbow trout *Oncorhynchus mykiss* (Walbaum), 359, 563
 rainbow trout, 145, 491, 891, 925
 rearing water, 747

- red blood cells, 165
 reproduction, 299, 1243
 reproductive
 performance, 129
 quality, 1095
 requirements, 395
 resource use efficiency, 403
 respiration, 33
 retention, 395
Rhombosolea tapirina, 425
 rice-fish culture, 969
rigor mortis, 703, 1059
 rohu, 547
 rotifers, 949

Saccostrea glomerata, 425
 salinity, 299, 425, 1009
 adaption, 891
Salmo salar L., 43, 425, 525, 533, 739, 875
 salmon, 533, 739, 875
 farming, 525
 salmonid, 799
 salmonidae, 563
 scallop, 819, 847
Scophthalmus maximus (Rafinesque), 177
 sea bass, 458, 569, 607, 1175
 sea cages, 501
Sebastes schlegelii (Hilgendorf), 979
 secondary responses, 777
 second-order polynomial regression, 673
 selectivity, 949
 self-feeding, 715
 behaviour, 509
Seriola quinqueradiata, 291
 sex categories, 1049
 sexual
 development, 863
 ratio, 607
 shell, 109
 density, 1019
 shape, 1019
 shrimp, 21, 231, 425, 785, 1073, 1129
 aquaculture, 785
 culture, 81
 effluent, 1
 farming, 241
 feed, 369
 feeding, 27, 995
 ponds, 269
 reproduction, 351
Sillago ciliata, 425
Silurus glanis, 999
 silver carp, 415
 silver perch, 1163
 simulation, 81
 size, 863, 959
 grading, 1221
 variation, 543
 slaughter, 37, 323, 643, 703
 smolt, 739
Solea senegalensis (Kaup), 747
 soybean meal, 279, 725
Sparus aurata (L. 1875), 747, 987
 spawn, 819

 spawning, 351, 863, 1243
 species selection, 95
 spectral composition, 251
 sperm, 1095
 spiny lobster, 913
 spotted babylon, 1265
 spotted wolffish *Anarhichas minor*, 1101, 1119
 stable isotopes, 839
 starvation, 197, 1090
 steam pelleting, 1163
 stickleback, 481
 stocking density, 621, 1031
 stress, 777, 829, 999
 stun, 37
 sturgeon, 725
 sunflower, 223
 Super Selco, 117
 supplementary feed, 151
 suppression, 445
 survival, 43, 637, 1119
 swimming behaviour, 501
 swordtail, 575

 T₁ treatment
 teleost fish, 383, 829
 temperature, 733, 959, 1073, 1119, 1175, 1243
 temperature/pH sex determination, 607
 tetra, 575
Tetraselmis, 949
 thermal
 biology, 525
 tolerance, 21
 tilapia, 223, 339, 941, 1199
 broodstock, 129
 tissue distribution, 1175
 tissue vitamin C, 261
 tolerance response, 747
 toxaphene, 925
 toxicants, 445
 toxicity, 1129
 transfer, 925
Trichoderma koningii, 1199
 triploid, 43
 trypsin, 333, 1073
 turbot (*Scophthalmus maximus* L.), 703
 turbot, 637

vibrio harveyi, 458
 Vietnam, 785

 water
 oxygenation, 607
 quality, 109, 151, 425, 785, 1103
 welfare, 37, 323
 wheat flour, 725
 white shrimp larvae, 333
 white shrimp, 995

 yabby *Cherax destructor* (Clark), 811, 913
 yellowtail flounder, 1193
 yellowtail, 291

 zeolite silicon, 75
 zeolites, 75

